

Transgenic manipulation of sn-glycerol-3-phosphate and glycerol production  
with a feedback defective glycerol-3-phosphate dehydrogenase gene

Ins.  
AI >

**Field of the invention**

5

The invention relates to the field of plant genetic engineering. More specifically, the invention relates to methods for manipulating the glycerol-3-phosphate metabolism of a plant by expressing in the plant a gene for a feedback defective glycerol-3-phosphate dehydrogenase.

10

**Background of the invention**

Glycerol-3-phosphate dehydrogenase (GPDH) (EC 1.1.1.8) is an essential enzyme for both prokaryotic and eukaryotic organisms. It catalyses the reduction of dihydroxyacetone phosphate (DHAP) to glycerol-3-phosphate (G-3-P) using NADH as reducing equivalent. Plant cells possess at least two isoforms of GPDH, one located in the plastids and the other in the cytosol<sup>1</sup>. The purification of the cytosolic GPDH from spinach has been reported<sup>2</sup>. The product of the reaction catalysed by GPDH, G-3-P, is a precursor for the synthesis of all glycerol lipid species, including membrane and storage lipids. The biosynthetic role of this enzyme in bacteria was established *in vivo* by the isolation of glycerol and G-3-P auxotrophs of *E. coli* mutant strains deficient in its activity<sup>3</sup>. These mutants could not synthesise phospholipid in the absence of supplemental G-3-P.

25

There are no reports of plant mutants defective in GPDH activity.

In addition to being essential for lipid biosynthesis, GPDH is involved in several other important biological processes. Most notably, GPDH, through consuming NADH and regenerating NAD<sup>+</sup>, plays an important role in maintaining cellular redox status. The NAD<sup>+</sup>/NADH couple plays a vital role

For catabolic reactions to proceed, the ratio  $\text{NAD}^+/\text{NADH}$  should be high.

GPDH reactions serves as a redox valve to dispose of extra reducing power. In this way, the cellular NAD<sup>+</sup>/NADH ratio can be maintained at a level allowing catabolic processes to proceed. The expression of the GPDH gene is subject to redox control and induced by anoxic conditions in *Saccharomyces cerevisiae*.

10 defective growth under anaerobic conditions<sup>4</sup>.

15 by a specific glycerol 3-phosphatase. To respond to a high external osmotic environment, yeast cells accumulate glycerol to compensate for differences between extracellular and intracellular water potentials<sup>5</sup>. The expression of the GPDH gene, *GPD1*, has been demonstrated to be osmoresponsive<sup>6</sup>. A strain of *Saccaromyces cerevisiae* in which the *GPD1* gene has been deleted is

hypersensitive to NaCl<sup>7</sup>. Accumulation of glycerol as an osmoregulatory solute has been reported in some halophilic green algae including *Dunaliella*, *Zooxanthellae*, *Asteromonas* and *Chlamydonas reinhardtii*<sup>8</sup>.

25 cytosolic isoform.

To date, there has been no report on the genetic manipulation of plant GPDH.

### Summary of the invention

It is an object of the invention to provide a method for expressing in a plant a heterologous glycerol-3-phosphate dehydrogenase.

- 5 It is an object of the invention to provide a plant expressing a heterologous glycerol-3-phosphate dehydrogenase, wherein the heterologous glycerol-3-phosphate dehydrogenase is subject to less feedback inhibition than wild type glycerol-3-phosphate dehydrogenase.

- 10 It is a further object of the invention to provide a genetically altered plant exhibiting altered fatty acid content in its glycerolipids.

It is a further object of the invention to provide a genetically altered plant exhibiting enhanced tolerance to osmotic stress in comparison to the wild type plant.

- 15 It is a further object of the invention to provide a genetically altered plant exhibiting increased stress tolerance in comparison to the wild type plant.

In a first aspect, the invention provides a method for expressing in a plant a heterologous glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition than wild type glycerol-3-phosphate dehydrogenase, the method comprising the steps of:

- 20 providing a vector comprising a DNA sequence encoding a glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition than wild type glycerol-3-phosphate dehydrogenase; and transforming the plant with the vector.

- 25 In a second aspect, the invention provides a plant expressing a heterologous glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition than wild type glycerol-3-phosphate dehydrogenase.

In a third aspect, the invention provides a method for producing a genetically altered plant having altered fatty acid content in its glycerolipids, the method comprising the steps of:

providing a vector comprising a DNA sequence encoding a  
glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition  
than wild type glycerol-3-phosphate dehydrogenase; and  
transforming the plant with the vector.

5 In a fourth aspect, the invention provides a method for producing a plant  
having increased glycerol and/or glycerol-3-phosphate levels, the method  
comprising the steps of:

providing a vector comprising a DNA sequence encoding a  
glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition  
10 than wild type glycerol-3-phosphate dehydrogenase; and  
transforming the plant with the vector.

In a fifth aspect, the invention provides a method for producing a  
genetically altered plant having increased stress tolerance relative to the wild  
type, the method comprising the steps of:

15 providing a vector comprising a DNA sequence encoding a  
glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition  
than wild type glycerol-3-phosphate dehydrogenase; and  
transforming the plant with the vector.

In a sixth aspect, the invention provides a method for producing a  
20 genetically altered plant having increased osmotic stress tolerance relative to  
the wild type, the method comprising the steps of:

providing a vector comprising a DNA sequence encoding a  
glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition  
than wild type glycerol-3-phosphate dehydrogenase; and  
25 transforming the plant with the vector.

In a seventh aspect, the invention provides a method for increasing the  
cellular glycerol-3-phosphate dehydrogenase activity in a plant, the method  
comprising the steps of:

providing a vector comprising a DNA sequence encoding a glycerol-3-phosphate dehydrogenase that is less sensitive to feedback inhibition than wild type glycerol-3-phosphate dehydrogenase; and transforming the plant with the vector.

- 5 In an eighth aspect, the invention provides a vector for genetically transforming a plant, wherein the vector comprises a DNA encoding a protein having glycerol-3-phosphate dehydrogenase activity, and the plant, after transforming, exhibits enhanced production of glycerol and/or glycerol-3-phosphate.

10

#### Detailed description of the invention

#### Brief description of the drawings

- 15 The invention is illustrated with the aid of the drawings, which show:  
FIG. 1 shows the nucleotide sequence and the deduced amino acid sequence of the *Escherichia coli* *gpsA2<sup>FR</sup>* gene. The point mutation is highlighted and denoted by '\*';

- FIG. 2 shows a diagram of the *gpsA2<sup>FR</sup>* plant transformation vector, pGPSA-VI, not drawn to scale;

- 20 FIG. 3 shows a southern blot analysis with respect to the *gpsA2<sup>FR</sup>* gene among the selected independent *Arabidopsis thaliana* transgenic lines.

FIG. 4 shows a northern blot analysis of *gpsA2<sup>FR</sup>* gene expression in the *A. thaliana* transgenic lines.

- 25 FIG. 5 shows the leaf fatty acid profiles of the selected *gpsA2<sup>FR</sup>* transgenic *Arabidopsis thaliana* lines.

FIG. 6 shows the germination rate of the seeds produced by the selected *Arabidopsis thaliana* transgenic lines in ½ MS medium with or without 225 mM NaCl.

FIG. 7 shows the germination rate of wild type *A. thaliana* and transgenic line #13 seeds in ½ MS media supplemented with various concentrations of NaCl.

FIG. 8 shows the performance of the soil-grow transgenic plants under various degree of salinity stress as detailed in Experimental Details.

Due to its role in lipid biosynthesis as well as in the stress responses, an increased GPDH activity in plants is desirable. Transgenic approaches to over express either a plant or a non-plant GPDH gene in a plant can, in principle, be expected to increase GPDH activity. However, there are several advantages inherent in inserting a non-plant gene into a plant genome. It is well established that introducing the same plant gene back to its originating species, even under sense-orientation, can result in a decrease of the over all enzyme activity due to co-suppression. Genes of different origin (heterologous), especially those from evolutionarily distantly related species, can be expected to be free of this impediment. More importantly, proteins of identical enzymatic function are often regulated through different schemes in different species. A heterologous enzyme may potentially be free of controlling factors that inhibit the endogenous enzyme.

The heterologous enzyme that is expressed in the plant, in the method of the invention, may be any glycerol-3-phosphate dehydrogenase that exhibits decreased inhibition of glycerol-3-phosphate production in the plant. Such enzymes are called feed-back defective. In a preferred embodiment, the heterologous enzyme is a glycerol-3-phosphate dehydrogenase having a single amino acid mutation. The mutation should not greatly decrease glycerol-3-phosphate dehydrogenase activity, but should decrease inhibition of the enzyme by glycerol-3-phosphate. One allele of the *E. coli* *gpsA* gene, *gpsA2<sup>FR</sup>*, has been reported to encode an altered version of the GPDH protein defective in feedback inhibition<sup>10</sup>. In a preferred embodiment, the method of the invention uses a vector comprising the gene *gpsA2<sup>FR</sup>*. The inventors identified a point mutation in the *gpsA2<sup>FR</sup>* sequence: replacement of A by C in

the third nucleotide of codon 255 in *gpsA*. The mutation results in substitution of Glu<sup>255</sup> (GAA) for Asp<sup>255</sup> (GAC) in the encoded protein. The sequences of the *gpsA*<sup>2FR</sup> gene and the deduced amino acid sequence of the gene are shown in FIG. 1. The gene sequence is listed in SEQ ID NO: 1, and the encoded protein is  
5 listed in SEQ ID NO: 2.

The vector may be any vector that is suitable for transforming the plant species used. Examples of suitable vectors include pHS737, pHS738, pRD400<sup>11</sup>; pBin19<sup>12</sup>; and pCGN3223<sup>13</sup>.

GPDH is common to the biosynthetic pathway of all plants. The method  
10 of the invention can therefore be used with any plant. The inventors chose to use the model plant species *Arabidopsis thaliana*. As a result of the ease with which this plant lends itself to work in both classical and molecular genetics, *Arabidopsis* has come to be widely used as a model organism in plant molecular genetics, development, physiology and biochemistry<sup>14,15,16</sup>. This  
15 dicotyledonous plant is also closely related to *Brassica* crop genus and it is increasingly apparent that information concerning the genetic control of basic biological processes in *Arabidopsis* will be transferable to other species<sup>17</sup>.

Indeed, there are numerous examples wherein studies of the molecular biology and biochemistry of a particular metabolic pathway or developmental  
20 process and the possibility of genetically engineering a plant to bring about changes to said metabolic pathway or process, has first been tested in the model plant *Arabidopsis*, and then shown to yield similar phenotypes in other plants, particularly crop plants.

Expressing a heterologous GPDH in a plant, according to the method of  
25 the invention, leads to altered fatty acid content in the triacylglycerols of the plant. It is often desirable to alter the fatty acid content of glycerolipids to achieve certain desired characteristics in oil seeds. For example, for oils destined for human consumption, it may be wished to increase unsaturated fatty acid content. For other uses, it may be desirable to increase the saturated  
30 fatty acid content. The inventors have found that plant transformants

over-expressing the *gpsA2<sup>FR</sup>* gene produce glycerolipids having an increased proportion of 16 carbon fatty acids and a concomitant decrease of 18 carbon fatty acids.

Due to the relationship of GPDH to glycerolipid synthesis, the method of the invention is particularly suited for use with oil seed bearing plants. The term oil seed bearing plant is meant to encompass any plant or crop from which the oil may be isolated in marketable quantity. Some plants or crops having glycerolipids with particularly interesting fatty acid composition are grown for the production of glycerolipids, even though the lipid content is low (e.g. less than 1 wt%). The method of the invention may be used in such plants to modify the fatty acid content of the glycerolipid. Preferred plants or crops are those having a seed lipid content of at least 1 wt%. Some illustrative examples of oil seed crops are as follows (trivial names are given in parentheses):

*Borago officinalis* (Borage); *Brassica* species, for example mustards, canola, rape, *B. campestris*, *B. napus*, *B. rapa*; *Cannabis sativa* (Hemp, widely uses as a vegetable oil in Asia); *Carthamus tinctorius* (Safflower); *Cocos nucifera* (Coconut); *Crambe abyssinica* (Crambe); *Cuphea* species (*Cuphea* produce medium chain fatty acids of industrial interest); *Elaeis guinensis* (African oil palm); *Elaeis oleifera* (American oil palm); *Glycine max* (Soybean); *Gossypium hirsutum* (Cotton - American); *Gossypium barbadense* (Cotton - Egyptian); *Gossypium herbaceum* (Cotton - Asiatic); *Helianthus annuus* (Sunflower); *Linum usitatissimum* (Linseed or flax); *Oenothera biennis* (Evening primrose); *Olea europea* (Olive); *Oryza sativa* (Rice); *Ricinus communis* (Castor); *Sesamum indicum* (Sesame); *Soja max* (Soybean - note *Glycine max* is the major species); *Triticum* species (Wheat); and *Zea maize* (Corn).

GPDH consumes NADH, and therefore plays an important role in maintaining a healthy cellular redox balance. Stress conditions often result in perturbation of plant metabolism, and particularly redox status. Stress conditions include such things as dryness, excessive humidity, excessive heat, excessive cold, excessive sunlight, and physical damage to the plant. Such



agents can lead to higher than normal levels of NADH. Excessive NADH can generate high concentrations of reactive oxygen species (ROS) that are hazardous to proteins and nucleic acids, and may even lead to cell death. An increased GPDH activity, as induced by the method of the invention, improves the capacity of plants to maintain cellular redox balance, thereby leading to an enhanced tolerance to stress.

Another type of stress suffered by plants is osmotic stress. This results when the plant is forced to grow in an environment in which the external water supply has an unusually high concentration of solute. The most usual solutes that are encountered include salts (particularly NaCl), however, in polluted areas, other solutes might be encountered. the method of the invention leads to increased levels of glycerol and/or glycerol-3-phosphate in the tissues of the transformed plant. Glycerol acts as an osmo-protectant, allowing the transformed plant to grow in conditions that would normally not support it.

A heterologous gene encoding GPDH activity can be introduced into genome of plants and expressed using conventional genetic engineering techniques. The most developed methodology for inserting genes into plant genomes is *Agrobacterium tumefaciens* mediated transformation. Other techniques known in the art of introducing DNA into plants include electroporation, chemically-mediated DNA uptake, and the use of microprojectiles.

The invention will be described in more detail with reference to the following examples. The examples serve only to illustrate the invention.

### Specific embodiments

#### a. Molecular Biological Techniques

For a general description of some of the techniques used, see Ausubel *et al* *Current protocols in Molecular Biology*, Vols 1, 2, 3, (1995) New York: Wiley, incorporated herein by reference.

b. Identification of the point mutation of the *gpsA2<sup>FR</sup>* gene from *Escherichia coli* strain BB26R.

In order to investigate the structure of the *gpsA2<sup>FR</sup>* gene, the inventors synthesised two primers, TTAGTGGCTGCTGCGCTC (GPSA3, SEQ ID NO: 3) and AACAAATGAACCAACGTAA (GPSA5, SEQ ID NO: 4), complementary to the sequences corresponding to the 3' and 5' end of the *gpsA* gene, respectively. PCR amplifications were performed with template DNA isolated from wild type *E. coli* K12 and from strain BB26R, respectively. The BB26R strain harbouring the *gpsA2<sup>FR</sup>* allele can be obtained according to Cronan *et al.*. The PCR products were purified with QIAquick™ PCR purification Kit (Qiagen™) and fully sequenced. The sequences of *gpsA* (wild type) and *gpsA2<sup>FR</sup>* (mutant) were compared through sequence alignment using the computer program DNASTar™.

c. Construction of a plant transformation vector for *gpsA<sup>FR</sup>*

Primers GAGAGCTCTTAGTGGCTGCTGCGCTC (GPSA31, SEQ ID NO: 5) and GAAGAAGGATCCAACAATGAACCAACGTAA (GPSA51, SEQ ID NO: 6) were designed according to the sequence of *gpsA2<sup>FR</sup>*. At the 5' end of GPSA31, a *SacI* restriction site was added, while a *BamHI* restriction site was added at the 5' end of GPSA5. The primers were used to perform PCR amplification of the *gpsA2<sup>FR</sup>* sequence. The PCR products were purified with QIAquick™ PCR purification Kit (Qiagen) and digested with *SacI/BamHI*. The *SacI/BamHI* digested *gpsA2<sup>FR</sup>* DNA fragment was subsequently inserted into the *Agrobacterium* binary vector pBI121 (Clontech) to replace the *SacI/BamHI* region covering the GUS gene. The resultant plant transformation vector is designated as pGPSA-VI (deposited August 31, 2000, at the American Type Culture Collection, 10801 University Blvd. Manassa, VA 20110-2209, accession no. PTA-2433). The *gpsA2<sup>FR</sup>* gene expression cassette in pGPSA-VI contains the *gpsA2<sup>FR</sup>*-encoding region driven by the constitutive 35S promoter. Its 3' end is flanked by the NOS terminator. The junction region between the 35S promoter

and the *gpsA2<sup>FR</sup>* encoding sequence in pGPSA-VI was confirmed through sequencing. The *gpsA2<sup>FR</sup>* protein will thus be expressed in all plant tissues including vegetative and reproductive (seed) tissues once the gene expression cassette is incorporated into the plant genome.

5

#### d. Plant Growth Conditions

*Arabidopsis thaliana* was chosen as the plant host to test the effect of the *gpsA2<sup>FR</sup>* gene since it is widely recognised as a laboratory model plant for genetic and biochemical studies. Moreover, *A. thaliana* in many aspects resembles *Brassica napus*, and is considered an oilseed plant. Genetic manipulations that are successful with *A. thaliana* can be applied to other species<sup>18</sup>. All *A. thaliana* control and transgenic plants were grown at the same time, in controlled growth chambers, under 16 hr fluorescent illumination (150-200  $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{sec}^{-1}$ ), 8 hr dark at 22 °C., as described previously<sup>19</sup>.

15

#### e. Plant Transformation

Plasmid pGPSA-VI was introduced into *Agrobacterium tumefaciens* strain GV3101 bearing helper nopaline plasmid pMP90, via electroporation. Wild type *A. thaliana* plants of ecotype Columbia were grown in soil. Plants one week after bolting were vacuum-infiltrated over night with a suspension of *A. tumefaciens* strain GV3101 harbouring pGPSA-VI <sup>20</sup>.

After infiltration, plants were grown to set seeds (T1). Dry seeds (T1) were harvested in bulk and screened on selective medium with 50 mg/L kanamycin. After two to three weeks on selective medium, kanamycin resistant seedlings (T1) which appeared as green were transformed to soil to allow growing to maturity. Seeds (T2) from the T1 plants were harvested and germinated on kanamycin plates to test segregation ratios. A typical single gene insertion event would give rise to a kanamycin resistant/sensitive ratio of 3:1. To further confirm the integration of the *gpsA2<sup>FR</sup>* gene, DNA was isolated from selected transgenic lines to perform Southern blot analysis with probes prepared with

30

- 25 Seed germination assays were performed with surface sterilised  
*Arabidopsis* seeds of wild type and selected T3 transgenic lines sown in Petri  
 dishes containing 20 ml half strength MS medium<sup>23</sup>, supplemented with B5  
 vitamins and 2% sucrose. For the salt stress germination assay, various  
 concentrations of NaCl were added. Cultures were grown at 22 °C under  
 30 fluorescent light, 16h light and 8h dark. Seed germination was recorded after a

period of 10 days. The emergence of radicle and cotyledons was considered as evidence of germination.

## Results

5

The *gpsA2<sup>FR</sup>* gene has a point mutation that alters one amino acid  
Residue in the GPDH protein (*gpsA2<sup>FR</sup>*)

The biosynthesis of G-3-P in *Escherichia coli* was initially investigated by Kito and Pizer<sup>24</sup>. The *gpsA* locus located at minute 71 of the *E. coli* genetic map  
10 was determined to be the structural gene for the biosynthetic glycerol-3-phosphate dehydrogenase by Cronan and Bell<sup>25</sup>. The nucleotide sequence and the deduced amino acid sequence of the *Escherichia coli* *gpsA* gene was reported previously<sup>26</sup>. Biochemical studies on phospholipid biosynthesis mutants indicated that the cellular level of G-3-P must be tightly regulated Bell (1974), *J.*  
15 *Bacteriol.* 117, 1065-1076]. The *E. coli* mutant, *plsB*, possesses a glycerol-P acyltransferase with an apparent  $K_m$  for G-3-P over 10 times higher than normal. Subsequently, revertants of the *plsB* mutant, BB26R, were identified<sup>27</sup>. The glycerol-3-phosphate dehydrogenase activities of these revertants were about 20-fold less sensitive to feedback inhibition by G-3-P. These feedback resistant  
20 *gpsA* alleles were named *gpsA2<sup>FR</sup>*. The molecular mechanism behind the *gpsA2<sup>FR</sup>* protein was unknown. The *gpsA2<sup>FR</sup>* gene was cloned from strain BB26R and its nucleotide sequence was determined. Sequence analysis indicated that *gpsA2<sup>FR</sup>* differs from *gpsA* at only one nucleotide base. The point mutation, a replacement of A from C at the third nucleotide of codon 255 in *gpsA*  
25 (FIG. 1) was founded in the *gpsA2<sup>FR</sup>* gene. This point mutation resulted in a change of Glu<sup>255</sup> (GAA) from Asp<sup>255</sup> (GAC) in the glycerol-3-phosphate dehydrogenase enzyme protein.

It has now been shown that the *gpsA2<sup>FR</sup>* gene harbours a point mutation in comparison to the wild type *gpsA* gene. The inventors have demonstrated that  
30 the point mutation is the reason why the GPDH enzyme is 20 time less sensitive

to G-3-P feedback inhibition than the wild type. As a result, the cellular G-3-P could reach a level higher than a wild type *gpsA* could generate.

**Introduction of the *gpsA2<sup>FR</sup>* gene into plant genomes does not affect  
5 plant development**

A large number of *gpsA2<sup>FR</sup>* transgenic plants were generated. These transgenic plants (T1) were initially screened for kanamycin resistance in kanamycin supplemented ½ MS medium. All T1 transgenic plants under our growing conditions appeared indistinguishable from wild type *A. thaliana*  
10 control, and developed at the same pace as that of the wild type plants when transferred into soil. The fertility and the seed yield were also not affected by the transgene. It thus proved that the integration of the *gpsA2<sup>FR</sup>* gene did not have any adversary effect on plant growth and reproduction. The segregation ratios of the (T2) seeds from the T1 plants with regard to kanamycin resistance  
15 were investigated. Transgenic line #7, #13, #54, #58 were selected for further study since segregation analysis indicated that these lines were single-insertion transgenic lines. To further verify the incorporation of *gpsA<sup>FR</sup>* gene into plant genome, genomic DNA was isolated from T3 plant seedlings of line #7, #13, #54, #58, respectively. Southern analysis of genomic DNA digested with three  
20 different restriction enzymes showed that these lines contain a single copy of the *gpsA2<sup>FR</sup>* gene, and the transgene is inherently stable (FIG. 4). Northern analysis with RNA extracted from these lines confirmed that the *gpsA2<sup>FR</sup>* gene is expressed at a high level in these transgenic lines. Therefore, the introduction and expression of the *gpsA2<sup>FR</sup>* gene into higher plants was accomplished.

25

***A. thaliana* *gpsA2<sup>FR</sup>* transformants have altered fatty acid profiles**

Total lipids were extracted from leaf tissues of transgenic plants as well as wild type control, and the fatty acid compositions were analysed using Gas Chromatography. In order to minimise any difference that might exist during  
30 plant development, care was taken to ensure all plant leaves collected were at

The present study revealed that the *gpsA2<sup>FR</sup>* transgenic plants possessed enhanced salinity tolerance.

Wild type *A. thaliana* could germinate reasonably well (80%) on medium containing 175 mM NaCl. However, seedling growth and development were severely retarded. In contrast, the growth rate of the transgenic plants was substantially higher. After 6 weeks, wild type plants developed chlorosis on leaf tissues and eventually died, while under the same conditions the transgenic plants still maintain relatively healthy green leaves. Plants growing in soil were also investigated with respect to salinity tolerance. The inventors followed the treatment protocol reported by Apse *et al*<sup>29</sup>, designed to mimic field stress conditions. As shown in FIG. 8, the transgenic plants displayed advanced growth and developmental profiles in comparison to those of wild type plants. Most of the wild type plants repeatedly treated with 50 mM NaCl appeared severely stressed with darkened leaf colour. The same treatment did not seem to affect the growth and reproduction of the transgenic lines. Wild type plants ceased to grow and eventually died when solutions containing salt at 100 mM were applied, while the majority of the transgenic plants developed to maturity and produced seeds. When a watering regime was carried out to a salt concentration of 150 mM NaCl, the transgenic plants showed apparent stressed

Wild type *A. thaliana* could germinate reasonably well (80%) on medium containing 175 mM NaCl. However, seedling growth and development were severely retarded. In contrast, the growth rate of the transgenic plants was substantially higher. After 6 weeks, wild type plants developed chlorosis on leaf tissues and eventually died, while under the same conditions the transgenic plants still maintain relatively healthy green leaves. Plants growing in soil were also investigated with respect to salinity tolerance. The inventors followed the treatment protocol reported by Apse *et al*<sup>29</sup>, designed to mimic field stress conditions. As shown in FIG. 8, the transgenic plants displayed advanced growth and developmental profiles in comparison to those of wild type plants. Most of the wild type plants repeatedly treated with 50 mM NaCl appeared severely stressed with darkened leaf colour. The same treatment did not seem to affect the growth and reproduction of the transgenic lines. Wild type plants ceased to grow and eventually died when solutions containing salt at 100 mM were applied, while the majority of the transgenic plants developed to maturity and produced seeds. When a watering regime was carried out to a salt concentration of 150 mM NaCl, the transgenic plants showed apparent stressed



phenotype, but were still able to produce seeds, albeit with short siliques and very little seed yield. Plants from line # 54 exhibited the most improved salinity among the transgenic lines tested. They produced seeds even when watering reached a salt concentration of 200 mM NaCl.

  
Tanya Nunnally, Patent Specialist  
ATCC Patent Depository

---

REFERENCES (incorporated herein by reference)

- <sup>1</sup> Gee *et al.*, (1988) *Plant Physiol.* 86, 98-103; Gee *et al.*, (1988) *Plant Physiol.* 87, 379-383.
- <sup>2</sup> Kirsh *et al.*, (1992) *Plant Physiol.* 100, 352-359
- <sup>3</sup> Hsu and Fox (1970) *J. Bacteriol.* 103, 410-416; Bell (1974) *J. Bacteriol.* 117, 1065-1076
- <sup>4</sup> Ansell *et al.*, (1997), *EMBO J.* 16, 2179-2187.
- <sup>5</sup> Brown (1990), in *Micorbial Water Stress Physiology, Principles and Perspectives*. John Wiley & Sons, New York
- <sup>6</sup> Larsson *et al.*, (1993), *Mol. Microbiol.* 10, 1101-1111.
- <sup>7</sup> Ansell *et al.*, (1997), *EMBO J.* 16, 2179-2187.
- <sup>8</sup> Husic and Tolbert, (1986), *Plant Physiol.* 82, 594-596; Ben-Amotz and Avron, (1983), *Annu Rev Microbiol.* 37, 95-119.
- <sup>9</sup> Hausmann *et al.*, (1995). In *Plant Lipid Metabolism*, (Kader, J.C., and Mazliak, P., eds), pp53-536, Kluwer Academic Publishers.
- <sup>10</sup> Bell and Cronan (1975), *J. Biol. Chem.* 250, 7147-7152.
- <sup>11</sup> Datla RS, Hammerlindl JK, Panchuk B, Pelcher LE, Keller W. (1992). *Modified binary plant transformation vectors with the wild-type gene encoding NPTII*; *Gene* 122:383-384.
- <sup>12</sup> Frisch DA, Harris-Haller LW, Yokubaitis NT, Thomas TL, Hardin SH, Hall TC. (1995). *Complete sequence of the binary vector Bin19*; *Plant Mol Biol* 27:405-409.
- <sup>13</sup> Roesler K, Shintani D, Savage L, Boddupalli S, Ohlrogge JB (1997) *Targeting of the Arabidopsis homomeric acetyl-coenzyme A carboxylase to plastids of rapeseeds*; *Plant Physiol* 113: 75-81
- <sup>14</sup> Meyerowitz, E.M. and Chang, C. (1985) Molecular biology of plant growth and development: *Arabidopsis thaliana* as an experimental system. In: *Developmental Biology*, Vol. 5, Plenum Press, NY., pp. 353-366.
- <sup>15</sup> Meyerowitz, E.M. (1987) *Arabidopsis thaliana*. *Ann. Rev. Genet.* 21: 93-111.
- <sup>16</sup> Goodman, H.M., Ecker, J.R. and Dean, C. (1995) The genome of *Arabidopsis thaliana*. *Proc. Nat'l. Acad. Sci. USA* 92: 10831-10835.
- <sup>17</sup> Lagercrantz, U., Putterill, J., Coupland, G. and Lydiate, D. (1996) Comparative mapping in *Arabidopsis* and *Brassica*, fine scale genome collinearity and congruence of genes controlling flowering. *Plant J.* 9: 13-20.
- <sup>18</sup> see for example: Zou *et al.*, United States Patent No: 6,051,755, April 18, 2000.
- <sup>19</sup> Katavic *et al.* (1995), *Plant Physiol.* 108:399-409.
- <sup>20</sup> Bechtolds *et al.* (C.R. Acad. Sci. Paris, *Sciences de la vie/Life sciences* 316, 1194-1199.
- <sup>21</sup> Katavic *et al.* (1995) *Plant Physiol.* 108:399-409.

- 
- <sup>22</sup> Apse *et al* (1999) *Science* 285, 1256-1258.
  - <sup>23</sup> Murashige and Skoog (1962), *Physiol Plant* 15: 473-497.
  - <sup>24</sup> *J. Biol. Chem.* (1969), 244, 3316-3333.
  - <sup>25</sup> Cronan and Bell; (1974), *J. Bacteriol.* 118, 598-605
  - <sup>26</sup> Ye and Larson (1988), *J. Bacteriol.*, 170, 4209-4215.
  - <sup>27</sup> Bell and Cronan (1975), *J. Biol. Chem.* 250, 7147-7152.
  - <sup>28</sup> Gardiner *et al* (1982), *Plant Physiol.* 70, 1316-1320.
  - <sup>29</sup> Apse *et al* (1999) *Science* 285, 1256-1258.